

Bayesian analysis of the astrobiological implications of life's early emergence on Earth

David S. Spiegel^{a,b,1} and Edwin L. Turner^{b,c}

^aSchool of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540; ^bDepartment of Astrophysical Sciences, Princeton University, Princeton, NJ 08544; and ^cInstitute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa 227-8568, Japan

Edited by Neta A. Bahcall, Princeton University, Princeton, NJ, and approved October 26, 2011 (received for review July 19, 2011)

Life arose on Earth sometime in the first few hundred million years after the young planet had cooled to the point that it could support water-based organisms on its surface. The early emergence of life on Earth has been taken as evidence that the probability of abiogenesis is high, if starting from young Earth-like conditions. We revisit this argument quantitatively in a Bayesian statistical framework. By constructing a simple model of the probability of abiogenesis, we calculate a Bayesian estimate of its posterior probability, given the data that life emerged fairly early in Earth's history and that, billions of years later, curious creatures noted this fact and considered its implications. We find that, given only this very limited empirical information, the choice of Bayesian prior for the abiogenesis probability parameter has a dominant influence on the computed posterior probability. Although terrestrial life's early emergence provides evidence that life might be abundant in the universe if early-Earth-like conditions are common, the evidence is inconclusive and indeed is consistent with an arbitrarily low intrinsic probability of abiogenesis for plausible uninformative priors. Finding a single case of life arising independently of our lineage (on Earth, elsewhere in the solar system, or on an extrasolar planet) would provide much stronger evidence that abiogenesis is not extremely rare in the universe.

Astrobiology is fundamentally concerned with whether extraterrestrial life exists and, if so, how abundant it is in the universe. The most direct and promising approach to answering these questions is surely empirical, the search for life on other bodies in the solar system (1, 2) and beyond in other planetary systems (3, 4). Nevertheless, a theoretical approach is possible in principle and could provide a useful complement to the more direct lines of investigation.

In particular, if we knew the probability per unit time and per unit volume of abiogenesis in a prebiotic environment as a function of its physical and chemical conditions and if we could determine or estimate the prevalence of such environments in the universe, we could make a statistical estimate of the abundance of extraterrestrial life. This relatively straightforward approach is, of course, thwarted by our great ignorance regarding both inputs to the argument at present.

There does, however, appear to be one possible way of finessing our lack of detailed knowledge concerning both the process of abiogenesis and the occurrence of suitable prebiotic environments (whatever they might be) in the universe. Namely, we can try to use our knowledge that life arose at least once in an environment (whatever it was) on the early Earth to try to infer something about the probability per unit time of abiogenesis on an Earth-like planet without the need (or ability) to say how Earth-like it need be or in what ways. We will hereinafter refer to this probability per unit time, which can also be considered a rate, as λ or simply the probability of abiogenesis.

Any inferences about the probability of life arising (given the conditions present on the early Earth) must be informed by how long it took for the first living creatures to evolve. By definition, improbable events generally happen infrequently. It follows that the duration between events provides a metric (however imperfect) of the probability or rate of the events. The time span be-

tween when Earth achieved prebiotic conditions suitable for abiogenesis plus generally habitable climatic conditions (5–7) and when life first arose, therefore, seems to serve as a basis for estimating λ . Revisiting and quantifying this analysis is the subject of this paper.

We note several previous quantitative attempts to address this issue in the literature, of which one (8) found, as we do, that early abiogenesis is consistent with life being rare, and the other (9) found that Earth's early abiogenesis points strongly to life being common on Earth-like planets (we compare our approach to the problem to that of ref. 9 below, including our significantly different results).^{*} Furthermore, an argument of this general sort has been widely used in a qualitative and even intuitive way to conclude that λ is unlikely to be extremely small because it would then be surprising for abiogenesis to have occurred as quickly as it did on Earth (10–16). Indeed, the early emergence of life on Earth is often taken as significant supporting evidence for optimism about the existence of extraterrestrial life (i.e., for the view that it is fairly common) (9, 17, 18). The major motivation of this paper is to determine the quantitative validity of this inference. We emphasize that our goal is not to derive an optimum estimate of λ based on all of the many lines of available evidence, but simply to evaluate the implication of life's early emergence on Earth for the value of λ .

A Bayesian Formulation of the Calculation

Bayes's theorem (19) can be written as $P[\mathcal{M}|\mathcal{D}] = (P[\mathcal{D}|\mathcal{M}] \times P_{\text{prior}}[\mathcal{M}]) / P[\mathcal{D}]$. Here, we take \mathcal{M} to be a model and \mathcal{D} to be data. To use this equation to evaluate the posterior probability of abiogenesis, we must specify appropriate \mathcal{M} and \mathcal{D} .

A Poisson or Uniform Rate Model. In considering the development of life on a planet, we suggest that a reasonable, if simplistic, model is that it is a Poisson process during a period of time from t_{min} until t_{max} . In this model, the conditions on a young planet preclude the development of life for a time period of t_{min} after its formation. Furthermore, if the planet remains lifeless until t_{max} has elapsed, it will remain lifeless thereafter as well because conditions no longer permit life to arise. For a planet around a solar-type star, t_{max} is almost certainly $\lesssim 10$ Gyr (10 billion years, the main sequence lifetime of the sun) and could easily be a substantially shorter period of time if there is something about the conditions on a young planet that are necessary for abiogenesis. Between these limiting times, we posit that there is a certain prob-

Author contributions: D.S.S. and E.L.T. designed research, performed research, analyzed data, and wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

^{*}There are two unpublished works (<http://arxiv.org/abs/0807.4969> and <http://www.vixra.org/abs/1108.0003>), of which we became aware after submission of this paper, that also conclude that early life on Earth does not rule out the possibility that abiogenesis is improbable.

¹To whom correspondence should be addressed. E-mail: dave@ias.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1111694108/-DCSupplemental.

ability per unit time (λ) of life developing. For $t_{\min} < t < t_{\max}$, then, the probability of life arising n times in time t is

$$P[\lambda, n, t] = P_{\text{Poisson}}[\lambda, n, t] = e^{-\lambda(t-t_{\min})} \frac{\{\lambda(t-t_{\min})\}^n}{n!}, \quad [1]$$

where t is the time since the formation of the planet.

This formulation could well be questioned on a number of grounds. Perhaps most fundamentally, it treats abiogenesis as though it were a single instantaneous event and implicitly assumes that it can occur in only a single way (i.e., by only a single process or mechanism) and only in one type of physical environment. It is, of course, far more plausible that abiogenesis is actually the result of a complex chain of events that take place over some substantial period of time and perhaps via different pathways and in different environments. However, knowledge of the actual origin of life on Earth, to say nothing of other possible ways in which it might originate, is so limited that a more complex model is not yet justified. In essence, the simple Poisson event model used in this paper attempts to integrate out all such details and treat abiogenesis as a black box process: Certain chemical and physical conditions as input produce a certain probability of life emerging as an output. Another issue is that λ , the probability per unit time, could itself be a function of time. In fact, the claim that life could not have arisen outside the window (t_{\min}, t_{\max}) is tantamount to saying that $\lambda = 0$ for $t \leq t_{\min}$ and for $t \geq t_{\max}$. Instead of switching from 0 to a fixed value instantaneously, λ could exhibit a complicated variation with time. If so, however, $P[\lambda, n, t]$ is not represented by the Poisson distribution and Eq. 1 is not valid. Unless a particular (non top-hat-function) time-variation of λ is suggested on theoretical grounds, it seems unwise to add such unconstrained complexity.

A further criticism is that λ could be a function of n : It could be that life arising once (or more) changes the probability per unit time of life arising again. Because we are primarily interested in the probability of life arising at all—i.e., the probability of $n \neq 0$ —we can define λ simply to be the value appropriate for a prebiotic planet (whatever that value may be) and remain agnostic as to whether it differs for $n \geq 1$. Thus, within the adopted model, the probability of life arising is one minus the probability of it not arising:

$$P_{\text{life}} = 1 - P_{\text{Poisson}}[\lambda, 0, t] = 1 - e^{-\lambda(t-t_{\min})}. \quad [2]$$

A Minimum Evolutionary Time Constraint. Naively, the single datum informing our calculation of the posterior of λ appears to be simply that life arose on Earth at least once, approximately 3.8 billion years ago (give or take a few hundred million years). There is additional significant context for this datum, however. Recall that the standard claim is that, because life arose early on the only habitable planet that we have examined for inhabitants, the probability of abiogenesis is probably high (in our language, λ is probably large). This standard argument neglects a potentially important selection effect; namely, on Earth, it took nearly 4 Gyr for evolution to lead to organisms capable of pondering the probability of life elsewhere in the universe. If this duration is necessary, then it would be impossible for us to find ourselves on, for example, a (~4.5-Gyr old) planet on which life first arose only after the passage of 3.5 billion years (20). On such planets there would not yet have been enough time for creatures capable of such contemplations to evolve. In other words, if evolution requires 3.5 Gyr for life to evolve from the simplest forms to intelligent, questioning beings, then we had to find ourselves on a planet where life arose relatively early, regardless of the value of λ (Table 1).

To introduce this constraint into the calculation we define δt_{evolve} as the minimum amount of time required after the emergence of life for cosmologically curious creatures to evolve, t_{emerge} as the age of the Earth from when the earliest extant evidence of

Table 1. Models of $t_0 = 4.5$ Gyr old planets

Model	Hypothetical	Conservative 1	Conservative 2	Optimistic
t_{\min}	0.5	0.5	0.5	0.5
t_{emerge}	0.51	1.3	1.3	0.7
t_{\max}	10	1.4	10	10
δt_{evolve}	1	2	3.1	1
t_{required}	3.5	1.4	1.4	3.5
Δt_1	0.01	0.80	0.80	0.20
Δt_2	3.00	0.90	0.90	3.00
\mathcal{R}	300	1.1	1.1	15

All times are in gigayears (Gyr). Two conservative models are shown, to indicate that t_{required} may be limited either by a small value of t_{\max} (Conservative 1), or by a large value of δt_{evolve} (Conservative 2).

life remains (though life might have actually emerged earlier), and t_0 as the current age of the Earth. The data, then, are that life arose on Earth at least once, approximately 3.8 billion years ago, and that this emergence was early enough that human beings had the opportunity subsequently to evolve and to wonder about their origins and the possibility of life elsewhere in the universe. In equation form, $t_{\text{emerge}} < t_0 - \delta t_{\text{evolve}}$.

The Likelihood Term. We now seek to evaluate the $P[\mathcal{D}|\mathcal{M}]$ term in Bayes's theorem. Let $t_{\text{required}} \equiv \min[t_0 - \delta t_{\text{evolve}}, t_{\max}]$. Our existence on Earth requires that life appeared within t_{required} . In other words, t_{required} is the maximum age that the Earth could have had at the origin of life in order for humanity to have a chance of showing up by the present. We define $\mathcal{S}_{\mathcal{E}}$ to be the set of all Earth-like worlds of age approximately t_0 in a large, unbiased volume and $L[t]$ to be the subset of $\mathcal{S}_{\mathcal{E}}$ on which life has emerged within a time t . $L[t_{\text{required}}]$ is the set of planets on which life emerged early enough that creatures curious about abiogenesis could have evolved before the present (t_0), and, presuming $t_{\text{emerge}} < t_{\text{required}}$ (which we know was the case for Earth), $L[t_{\text{emerge}}]$ is the subset of $L[t_{\text{required}}]$ on which life emerged as quickly as it did on Earth. Correspondingly, $N_{\mathcal{S}_{\mathcal{E}}}$, N_{t_0} , and N_{t_e} are the respective numbers of planets in sets $\mathcal{S}_{\mathcal{E}}$, $L[t_{\text{required}}]$, and $L[t_{\text{emerge}}]$. The fractions $\varphi_{t_0} \equiv N_{t_0}/N_{\mathcal{S}_{\mathcal{E}}}$ and $\varphi_{t_e} \equiv N_{t_e}/N_{\mathcal{S}_{\mathcal{E}}}$ are, respectively, the fraction of Earth-like planets on which life arose within t_{required} and the fraction on which life emerged within t_{emerge} . The ratio $r \equiv \varphi_{t_e}/\varphi_{t_0} = N_{t_e}/N_{t_0}$ is the fraction of L_{t_0} on which life arose as soon as it did on Earth. Given that we had to find ourselves on such a planet in the set L_{t_0} to write and read about this topic, the ratio r characterizes the probability of the data given the model if the probability of intelligent observers arising is independent of the time of abiogenesis (so long as abiogenesis occurs before t_{required}). (This last assumption might seem strange or unwarranted, but the effect of relaxing this assumption is to make it more likely that we would find ourselves on a planet with early abiogenesis and therefore to reduce our limited ability to infer anything about λ from our observations.) Because $\varphi_{t_e} = 1 - P_{\text{Poisson}}[\lambda, 0, t_{\text{emerge}}]$ and $\varphi_{t_0} = 1 - P_{\text{Poisson}}[\lambda, 0, t_{\text{required}}]$, we may write that

$$P[\mathcal{D}|\mathcal{M}] = \frac{1 - \exp[-\lambda(t_{\text{emerge}} - t_{\min})]}{1 - \exp[-\lambda(t_{\text{required}} - t_{\min})]}, \quad [3]$$

if $t_{\min} < t_{\text{emerge}} < t_{\text{required}}$ (and $P[\mathcal{D}|\mathcal{M}] = 0$ otherwise). This function is called the Likelihood function, and represents the probability of the observation(s), given a particular model.[†] It is via this function that the data condition our prior beliefs about λ in standard Bayesian terminology.

[†]An alternative way to derive Eq. 3 is to let E = abiogenesis occurred between t_{\min} and t_{emerge} and R = abiogenesis occurred between t_{\min} and t_{required} . We then have, from the rules of conditional probability, $P[E|R, \mathcal{M}] = P[E, R, \mathcal{M}]/P[R, \mathcal{M}]$. Because E entails R , the numerator on the right-hand side is simply equal to $P[E, \mathcal{M}]$, which means that the previous equation reduces to Eq. 3.

Limiting Behavior of the Likelihood. It is instructive to consider the behavior of Eq. 3 in some interesting limits. For $\lambda(t_{\text{required}} - t_{\text{min}}) \ll 1$, the numerator and denominator of Eq. 3 each go approximately as the argument of the exponential function; and therefore, in this limit, the Likelihood function is approximately constant:

$$P[\mathcal{D}|\mathcal{M}] \approx \frac{t_{\text{emerge}} - t_{\text{min}}}{t_{\text{required}} - t_{\text{min}}} \quad [4]$$

This result is intuitively easy to understand as follows: if λ is sufficiently small, it is overwhelmingly likely that abiogenesis occurred only once in the history of the Earth, and by the assumptions of our model, the one event is equally likely to occur at any time during the interval between t_{min} and t_{required} . The chance that this abiogenesis event will occur by t_{emerge} is then just the fraction of that total interval that has passed by t_{emerge} —the result given in Eq. 4.

In the other limit, when $\lambda(t_{\text{emerge}} - t_{\text{min}}) \gg 1$, the numerator and denominator of Eq. 3 are both approximately 1. In this case, the Likelihood function is also approximately constant (and equal to unity). This result is even more intuitively obvious because a very large value of λ implies that abiogenesis events occur at a high rate (given suitable conditions) and are thus likely to have occurred very early in the interval between t_{min} and t_{required} .

These two limiting cases, then, already reveal a key conclusion of our analysis: The posterior distribution of λ for both very large and very small values will have the shape of the prior, just scaled by different constants. Only when λ is neither very large nor very small—or, more precisely, when $\lambda(t_{\text{emerge}} - t_{\text{min}}) \approx 1$ —do the data and the prior both inform the posterior probability at a roughly equal level.

The Bayes Factor. In this context, note that the probability in Eq. 3 depends crucially on two time differences, $\Delta t_1 \equiv t_{\text{emerge}} - t_{\text{min}}$ and $\Delta t_2 \equiv t_{\text{required}} - t_{\text{min}}$, and that the ratio of the Likelihood function at large λ to its value at small λ goes roughly as

$$\mathcal{R} \equiv \frac{P[\text{data}|\text{large}\lambda]}{P[\text{data}|\text{small}\lambda]} \approx \frac{\Delta t_2}{\Delta t_1} \quad [5]$$

\mathcal{R} is called the Bayes factor or Bayes ratio and is sometimes employed for model selection purposes. In one conventional interpretation (21), $\mathcal{R} \leq 10$ implies no strong reason in the data alone to prefer the model in the numerator over the one in the denominator. For the problem at hand, this interpretation means that the datum does not justify preference for a large value of λ over an arbitrarily small one unless Eq. 5 gives a result larger than roughly ten.

Because the Likelihood function contains all of the information in the data and because the Bayes factor has the limiting behavior given in Eq. 5, our analysis in principle need not consider priors. If a small value of λ is to be decisively ruled out by the data, the value of \mathcal{R} must be much larger than unity. It is not for plausible choices of the parameters (see Table 1), and thus arbitrarily small values of λ can only be excluded by some adopted prior on its values. Still, for illustrative purposes, we now proceed to demonstrate the influence of various possible λ priors on the λ posterior.

The Prior Term. To compute the desired posterior probability, what remains to be specified is $P_{\text{prior}}[\mathcal{M}]$, the prior joint probability density function (PDF) of λ , t_{min} , t_{max} , and δt_{evolve} . One approach to choosing appropriate priors for t_{min} , t_{max} , and δt_{evolve} , would be to try to distill geophysical and paleobiological evidence along with theories for the evolution of intelligence and the origin of life into quantitative distribution functions that accurately represent prior information and beliefs about these parameters. Then,

to ultimately calculate a posterior distribution of λ , one would marginalize over these nuisance parameters. However, because our goal is to evaluate the influence of life's early emergence on our posterior judgment of λ (and not of the other parameters), we instead adopt a different approach. Rather than calculating a posterior over this four-dimensional parameter space, we investigate the way these three time parameters affect our inferences regarding λ by simply taking their priors to be delta functions at several theoretically interesting values: a purely hypothetical situation in which life arose extremely quickly, a most conservative situation, and an in between case that is also optimistic but for which there does exist some evidence (see Table 1).

For the values in Table 1, the likelihood ratio \mathcal{R} varies from approximately 1.1 to 300, with the parameters of the optimistic model giving a borderline significance value of $\mathcal{R} = 15$. Thus, only the hypothetical case gives a decisive preference for large λ by the Bayes factor metric, and we emphasize that there is no direct evidence that abiogenesis on Earth occurred that early, only 10 million years after conditions first permitted it.*

We also lack a first-principles theory or other solid prior information for λ . We therefore take three different functional forms for the prior—uniform in λ , uniform in λ^{-1} (equivalent to saying that the mean time until life appears is uniformly distributed), and uniform in $\log_{10} \lambda$. For the uniform in λ prior, we take our prior confidence in λ to be uniformly distributed on the interval 0 to $\lambda_{\text{max}} = 1,000 \text{ Gyr}^{-1}$ (and to be 0 otherwise). For the uniform in λ^{-1} and the uniform in $\log_{10}[\lambda]$ priors, we take the prior density functions for λ^{-1} and $\log_{10}[\lambda]$, respectively, to be uniform on $\lambda_{\text{min}} \leq \lambda \leq \lambda_{\text{max}}$ (and 0 otherwise). For illustrative purposes, we take three values of λ_{min} : $10^{-22} \text{ Gyr}^{-1}$, $10^{-11} \text{ Gyr}^{-1}$, and 10^{-3} Gyr^{-1} , corresponding roughly to life occurring once in the observable universe, once in our galaxy, and once per 200 stars (assuming one Earth-like planet per star).

In standard Bayesian terminology, both the uniform in λ and the uniform in λ^{-1} priors are said to be highly informative. This appellation means that they strongly favor large and small, respectively, values of λ in advance, i.e., on some basis other than the empirical evidence represented by the likelihood term. For example, the uniform in λ prior asserts that we know on some other basis (other than the early emergence of life on Earth) that it is a hundred times less likely that λ is less than 10^{-3} Gyr^{-1} than that it is less than 0.1 Gyr^{-1} . The uniform in λ^{-1} prior has the equivalent sort of preference for small λ values. By contrast, the logarithmic prior is relatively uninformative in standard Bayesian terminology and is equivalent to asserting that we have no prior information that informs us of even the order-of-magnitude of λ .

In our opinion, the logarithmic prior is the most appropriate one given our current lack of knowledge of the process(es) of abiogenesis, as it represents scale-invariant ignorance of the value of λ . It is, nevertheless, instructive to carry all three priors through the calculation of the posterior distribution of λ , because they vividly illuminate the extent to which the result depends on the data versus the assumed prior.

Comparison with Previous Analysis. Using a binomial probability analysis, Lineweaver and Davis (9) attempted to quantify q , the probability that life would arise within the first billion years on an Earth-like planet. Although the binomial distribution typically applies to discrete situations (in contrast to the continuous passage of time, during which life might arise), there is a simple correspondence between their analysis and the Poisson model described above. The probability that life would arise at least once within a billion years (what ref. 9 calls q) is a simple transformation of λ , obtained from Eq. 2, with $\Delta t_1 = 1 \text{ Gyr}$:

*Ref. 22 advances this claim based on theoretical arguments that are critically reevaluated in ref. 23.

$$q = 1 - e^{-(\lambda)(1 \text{ Gyr})} \quad \text{or} \quad \lambda = \ln[1 - q]/(1 \text{ Gyr}). \quad [6]$$

In the limit of $\lambda(1 \text{ Gyr}) \ll 1$, Eq. 6 implies that q is equal to $\lambda(1 \text{ Gyr})$. Though not cast in Bayesian terms, the analysis in ref. 9 draws a Bayesian conclusion and therefore is based on an implicit prior that is uniform in q . As a result, it is equivalent to our uniform- λ prior for small values of λ (or q), and it is this implicit prior, not the early emergence of life on Earth, that dominates their conclusions.

The Posterior Probability of Abiogenesis

We compute the normalized product of the probability of the data given λ (Eq. 3) with each of the three priors (uniform, logarithmic, and inverse-uniform). This computation gives us the Bayesian posterior PDF of λ , which we also derive for each model in Table 1. Then, by integrating each PDF from $-\infty$ to λ , we obtain the corresponding cumulative distribution function (CDF).

Fig. 1, displays the results by plotting the prior and posterior probability of λ . The *Left* presents the PDF, and the *Right* the CDF, for uniform, logarithmic, and inverse-uniform priors, for model optimistic, which sets Δt_1 (the maximum time it might have taken life to emerge once Earth became habitable) to 0.2 Gyr, and Δt_2 (the time life had available to emerge in order that intelligent creatures would have a chance to evolve) to 3.0 Gyr. The dashed and solid curves represent, respectively, prior and posterior probability functions. In this figure, the priors on λ have $\lambda_{\min} = 10^{-3} \text{ Gyr}^{-1}$ and $\lambda_{\max} = 10^3 \text{ Gyr}^{-1}$. The green, blue, and red curves are calculated for uniform, logarithmic, and inverse-uniform priors, respectively. The results of the corresponding calculations for the other models and bounds on the assumed priors are presented in *SI Text*, but the cases shown in Fig. 1 suffice to demonstrate all of the important qualitative behaviors of the posterior.

In the plot of differential probability (PDF, *Left*), it appears that the inferred posterior probabilities of different values of λ are conditioned similarly by the data (leading to a jump in the posterior PDF of roughly an order-of-magnitude in the vicinity

of $\lambda \sim 0.5 \text{ Gyr}^{-1}$). The plot of cumulative probability, however, immediately shows that the uniform and the inverse priors produce posterior CDFs that are completely insensitive to the data. Namely, small values of λ are strongly excluded in the uniform in λ prior case and large values are equally strongly excluded by the uniform in λ^{-1} prior, but these strong conclusions are not a consequence of the data, only of the assumed prior. This point is particularly salient, given that a Bayesian interpretation of ref. 9 indicates an implicit uniform prior. In other words, their conclusion that q cannot be too small and thus that life should not be too rare in the universe is not a consequence of the evidence of the early emergence of life on Earth but almost only of their particular parameterization of the problem.

For the optimistic parameters, the posterior CDF computed with the uninformative logarithmic prior does reflect the influence of the data, making greater values of λ more probable in accordance with one's intuitive expectations. However, with this relatively uninformative prior, there is a significant probability that λ is very small (12% chance that $\lambda < 1 \text{ Gyr}^{-1}$). Moreover, if we adopted smaller λ_{\min} , smaller λ_{\max} , and/or a larger $\Delta t_1/\Delta t_2$ ratio, the posterior probability of an arbitrarily low λ value can be made acceptably high (see Fig. 2 and *SI Text*).

Independent Abiogenesis. We have no strong evidence that life ever arose on Mars (although no strong evidence to the contrary either). Recent observations have tentatively suggested the presence of methane at the level of approximately 20 ppb (24), which could potentially be indicative of biological activity. The case is not entirely clear, however, as alternative analysis of the same data suggests that an upper limit to the methane abundance is in the vicinity of approximately 3 ppb (25). If, in the future, researchers find compelling evidence that Mars or an exoplanet hosts life that arose independently of life on Earth [or that life arose on Earth a second, independent time (26, 27)], how would this discovery affect the posterior probability density of λ (assuming that the same λ holds for both instances of abiogenesis)?




Fig. 1. PDF and CDF of λ for uniform, logarithmic, and inverse-uniform priors, for model optimistic, with $\lambda_{\min} = 10^{-3} \text{ Gyr}^{-1}$ and $\lambda_{\max} = 10^3 \text{ Gyr}^{-1}$. (*Left*) The dashed and solid curves represent, respectively, the prior and posterior PDFs of λ under three different assumptions about the nature of the prior. The green curves are for a prior that is uniform on the range $0 \text{ Gyr}^{-1} \leq \lambda \leq \lambda_{\max}$ (uniform); the blue are for a prior that is uniform in the log of λ on the range $-3 \leq \log \lambda \leq 3$ [Log (-3)]; and the red are for a prior that is uniform in λ^{-1} on the interval $10^{-3} \text{ Gyr} \leq \lambda^{-1} \leq 10^3 \text{ Gyr}$ [InvUnif (-3)]. (*Right*) The curves represent the CDFs of λ . The ordinate on each curve represents the integrated probability from 0 to the abscissa (color and line-style schemes are the same as in *Left*). For a uniform prior, the posterior CDF traces the prior almost exactly. In this case, the posterior judgment that λ is probably large simply reflects the prior judgment of the distribution of λ . For the prior that is uniform in λ^{-1} (InvUnif), the posterior judgment is quite opposite—namely, that λ is probably quite small—but this judgment is also foretold by the prior, which is traced nearly exactly by the posterior. For the logarithmic prior, the datum (that life on Earth arose within a certain time window) does influence the posterior assessment of λ , shifting it in the direction of making greater values of λ more probable. Nevertheless, the posterior probability is approximately 12% that $\lambda < 1 \text{ Gyr}^{-1}$. Lower λ_{\min} and/or lower λ_{\max} would further increase the posterior probability of very low λ , for any of the priors.

studies might, in the near future, help us to significantly refine our estimate of the probability (per unit time, per Earth-like planet) of abiogenesis.

ACKNOWLEDGMENTS. We thank Adam Brown, Adam Burrows, Chris Chyba, Scott Gaudi, Aaron Goldman, Alan Guth, Larry Guth, Laura Landweber, Tullis Onstott, Caleb Scharf, Stanley Spiegel, Josh Winn, and Neil Zimmerman for thoughtful discussions. E.L.T. is grateful to Carl Boettiger for calling this

problem to his attention some years ago. D.S.S. acknowledges support from National Aeronautics and Space Administration Grant NNX07AG80G and from Jet Propulsion Laboratory/Spitzer Agreements 1328092, 1348668, and 1312647. E.L.T. gratefully acknowledges support from the World Premier International Research Center Initiative (Institute for the Physics and Mathematics of the Universe), Ministry of Education, Culture, Sports, Science, and Technology, Japan, and the Research Center for the Early Universe at the University of Tokyo as well as the hospitality of its Department of Physics.

1. Chyba CF, Hand KP (2005) Astrobiology: The study of the living universe. *Ann Rev Astr Ap* 43:31–74.
2. Des Marais D, Walter M (1999) Astrobiology: Exploring the origins, evolution, and distribution of life in the universe. *Annu Rev Ecol Syst* 397–420.
3. Des Marais DJ, et al. (2002) Remote sensing of planetary properties and biosignatures on extrasolar terrestrial planets. *Astrobiology* 2:153–181.
4. Seager S, Turner EL, Schafer J, Ford EB (2005) Vegetation's red edge: A possible spectroscopic biosignature of extraterrestrial plants. *Astrobiology* 5:372–390.
5. Kasting JF, Whitmire DP, Reynolds RT (1993) Habitable zones around main sequence stars. *Icarus* 101:108–128.
6. Selsis F, et al. (2007) Habitable planets around the star Gliese 581? *Astron Astrophys* 476:1373–1387.
7. Spiegel DS, Menou K, Scharf CA (2008) Habitable climates. *Astrophys J* 681:1609–1623.
8. Carter B (1983) The anthropic principle and its implications for biological evolution. *Philos Trans R Soc A* 310:347–363.
9. Lineweaver CH, Davis TM (2002) Does the rapid appearance of life on Earth suggest that life is common in the universe? *Astrobiology* 2:293–304.
10. van Zuilen MA, Lepland A, Arrhenius G (2002) Reassessing the evidence for the earliest traces of life. *Nature* 418:627–630.
11. Westall F (2005) Early life on Earth: The ancient fossil record. *Astrobiology: Future Perspectives*, eds P Ehrenfreund et al. (Springer, Berlin), 305, pp 287–316.
12. Moorbath S (2005) Oldest rocks, earliest life, heaviest impacts, and the hadea archaean transition. *Appl Geochem* 20:819–824.
13. Nutman A, Friend C (2006) Petrography and geochemistry of apatites in banded iron formation, Akilia, W. Greenland: Consequences for oldest life evidence. *Precambrian Res* 147:100–106.
14. Buick R (2007) The earliest records of life on Earth. *Planets and Life: The Emerging Science of Astrobiology*, eds WT Sullivan and JA Baross (Cambridge University Press, New York).
15. Sullivan W, Baross J (2007) *Planets and Life: The Emerging Science of Astrobiology* (Cambridge University Press, New York).
16. Sugitani K, et al. (2010) Biogenicity of morphologically diverse carbonaceous microstructures from the ca. 3,400 Ma Strelley Pool Formation, in the Pilbara Craton, Western Australia. *Astrobiology* 10:899–920.
17. Ward PD, Brownlee D (2000) *Rare Earth: Why complex life is uncommon in the universe.* (Springer, New York), pp 66–67 p 243.
18. Darling D (2001) *Life Everywhere* (Basic Books).
19. Bayes M, Price M (1763) An essay toward solving a problem in the doctrine of chances. by the late Rev. Mr. Bayes, FRS communicated by Mr. Price, in a letter to John Canton, AMFRS. *Phil Trans* 53:370–418.
20. Lineweaver CH, Davis TM (2003) On the nonobservability of recent biogenesis. *Astrobiology* 3:241–243.
21. Jeffreys H (1961) *Theory of Probability* (Oxford Univ Press, Oxford).
22. Lazcano A, Miller S (1994) How long did it take for life to begin and evolve to cyanobacteria? *J Mol Evol* 39:546–554.
23. Orgel L (1998) The origin of life-how long did it take? *Origins Life Evol Biospheres* 28:91–96.
24. Mumma M, et al. (2009) Strong release of methane on Mars in northern summer 2003. *Science* 323:1041–1045.
25. Zahnle K, Freedman RS, Catling DC (2010) Is there methane on Mars? *Icarus*, 212 pp:493–503.
26. Davies PCW, Lineweaver CH (2005) Finding a second sample of life on Earth. *Astrobiology* 5:154–163.
27. Davies PCW, et al. (2009) Signatures of a shadow biosphere. *Astrobiology* 9:241–249.
28. Vogt SS, et al. (2010) The Lick-Carnegie exoplanet survey: A 3.1 earth-mass planet in the habitable zone of the nearby M3V star Gliese 581. *Astrophys J* 723:954–965.
29. Borucki WJ, et al. (2011) Characteristics of planetary candidates observed by Kepler, II: Analysis of the first four months of data. *Astrophys J* 736:19.
30. Howard AW, et al. (2011) Planet occurrence within 0.25 AU of solar-type stars from Kepler. arXiv:1103.2541v1.
31. Wordsworth RD, et al. (2011) Gliese 581d is the first discovered terrestrial-mass exoplanet in the habitable zone. *Astrophys J Lett* 733:L48.
32. Kaltenegger L, Selsis F (2007) Biomarkers set in context. ArXiv:0710.0881.
33. Tarter J (2001) The search for extraterrestrial intelligence (SETI). *Annu Rev Astron Astrophys* 39:511–548.